

COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

What are foetal alcohol spectrum disorders (FASD)?

FASD is the result of alcohol being drunk while pregnant. FASD includes foetal alcohol syndrome (FAS), which is the easiest to recognise, as the students have certain facial features. These can include a thin upper lip, a flat philtrum (the groove between the nose and upper lip) and smaller eye openings. Although students with FASD may need support with some behaviours, they also have strengths.

Some common **strengths** are:

- strong verbal skills
- being bright in some areas, such as being artistic, musical, athletic, or having practical skills
- a friendly, affectionate, helpful and generous nature
- being good with younger children
- determination and persistence.

Likely **difficulties** are:


- memory problems; difficulty storing and retrieving information
- inconsistent performance (on and off days)
- being impulsive, easily distracted and disorganised
- an ability to repeat instructions, but not to put them into action
- difficulty with abstract concepts such as maths, money management and time
- being slow to process information; needing time to think
- being slow to follow what is said to them, such as only taking in every third word
- developmental delays, which make them seem younger than their actual age
- inability to understand the consequences of their actions.

Strategies that may help

Students with FASD are affected in different ways and to different degrees, depending on what stage of the pregnancy alcohol was consumed. This means that every student's pattern of difficulties will be different to some extent and, unlike students who are slow to develop, these students may have parts of the brain that were never fully formed. This can make learning some things extremely hard. Often, maths is a particular area of difficulty.

The following strategies are useful in supporting a student with FASD.

- 1 Closely supervise the student to ensure their safety.
- 2 Keep to the same routines, and help students to follow the structure of the day.
- 3 Keep in close contact with family or carers so that there is a consistent approach.




COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

- 4 Keep instructions short and simple. Support what is said with visually presented material as well, such as visual timetables, symbols and pictures, etc.
- 5 Present tasks in a similar way each time, so that students know what to do.
- 6 Use students' strengths so that they can experience success. Help them to gain satisfaction from their work and celebrate their successes.
- 7 Teach them how to communicate their feelings; for instance, by talking about pictures of people showing different emotions. Help them to understand how they feel and to learn the words to describe their feelings.
- 8 Help them to become more organised, for instance, by having pictures or lists of the equipment they need for a lesson or activity. Keep the classroom as organised as possible, and label everything that they might need.
- 9 Help them to understand and remember school or class rules by discussing them regularly. It is helpful if these are positive statements rather than a list of 'Don't do's. If rules are broken, correct the behaviour immediately in a supportive way and practise the right behaviour.
- 10 Work on social skills and life skills as well as helping students with their work.

It is thought that up to 80% of children with FASD are fostered or adopted, so they need the school to provide as much support, stability and encouragement as possible.

It is important to remember that every student has their own likes and interests, and to use these to help them to learn. Families may also have ideas they would be happy to share, and may like to hear about what strategies work in the classroom. It will help the student if their family and the school work together.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

What is foetal alcohol spectrum disorder (FASD)?

FASD is an umbrella term used to encompass the range of possible effects of prenatal exposure to alcohol (British Medical Association, 2007). The following diagnostic or educational terms are included in the spectrum:

- **foetal alcohol syndrome (FAS)** – the most easily recognisable condition due to characteristic facial features which are formed during the first trimester of pregnancy and dissipate with age
- **partial foetal alcohol syndrome (pFAS)** – some but not all of the criteria for FAS are met
- **foetal alcohol effects (FAE)** – the symptoms are not usually visible (eg behaviour disorders, attention deficits, etc)
- **alcohol related neurodevelopmental disorder (ARND)** – can include attention deficits, behaviour disorders, obsessive/compulsive disorder
- **alcohol related birth defects (ARBD)** – includes characteristics such as organ damage, heart defects, sight/hearing problems, skeletal damage and joint defects

A diagnosis of FAS is not an indication of the severity of the impairment, which varies considerably. For a diagnosis, four criteria must be met:

- growth deficiency
- characteristic facial features
- central nervous system damage
- confirmed alcohol exposure.

The characteristics of students

The characteristics of FASD are individual to each student depending on timing and dosage of maternal alcohol consumption, and maternal well-being and health during pregnancy. Damage to the central nervous system results in changes to the structure of the brain that will persist throughout life. Permanent damage to organs and bones can occur at particular periods of pregnancy. Generally, however, the following characteristics may be observed in these students:

Possible strengths

Students may be:

- bright in some areas; highly verbal
- artistic, musical, athletic, have good practical skills
- friendly, helpful, affectionate and good with younger students
- determined and persistent.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

Likely difficulties

Students may:

- be easily influenced by others
- have difficulty predicting and understanding the consequences of actions
- despite a good vocabulary, struggle to understand what is said to them
- have difficulty in separating fact from fantasy
- display behaviours which will need support, such as lying, stealing, temper tantrums
- have delayed physical, emotional and cognitive development (delayed developmental milestones)
- have poor impulse control, hyperactivity and poor memory
- experience sensory processing difficulties
- have social communication difficulties, particularly in the areas of interpreting the actions of others and in understanding how to respond in social situations; this is due to their impulsivity and lack of inhibition, and may lead to inappropriate sexual behaviour when older
- have dietary and feeding difficulties/small stature/health problems such as frequent and persistent colds, visual and hearing impairments.


Implications for teaching and learning

As a result of their disability, students may experience:

- memory problems; difficulty storing and retrieving information
- inconsistent performance (on and off days)
- impulsivity, distractibility, disorganisation
- ability to repeat instructions, but inability to put them into action
- difficulty with abstractions, such as maths, money management, time concepts
- cognitive processing deficits, so need time to take in information and to respond
- slow auditory pace (may only understand every third word of normally paced conversation)
- developmental lags (may act younger than chronological age)
- inability to predict outcomes or understand consequences.

Supporting students with FASD

1. Have a safe, structured environment, where the student's difficulty in following verbal instruction is supported by visually presented material. If helpful, use screens or something similar to reduce the distraction around the area where they work. Keep instructions short and simple.
2. Support their independence and organisational skills by creating clearly defined areas of the classroom for specific activities; use labelling and general tidiness to help them to find what they need.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

3. Have a set routine for carrying out their work, so they know what is expected of them.
4. Allow them time to take in what is said and to form a response. Provide plenty of repetition to aid recall.
5. Check that they understand class and school rules and give them the chance to go over them regularly.
6. Provide plenty of encouragement and praise when they achieve. Use their strengths and interest to personalise learning and assist motivation.
7. Help them to understand their feelings and practice the vocabulary to express how they feel. (It is thought that up to 80% of children affected by FASD are with foster or adoptive families, and that some may have lived with a number of families before they are placed with adoptive families. These experiences may lead to impaired emotional understanding.)
8. Maintain regular contact with home in order to give consistent messages and to provide support.
9. The transition between primary and secondary education can be difficult for students with FASD, as they may already have had a number of changes in their lives. Try to ensure that there is a smooth transition and that support remains in place.
10. For teenagers, issues around emotions, friendships and sexual behaviour, independence and achievement can be difficult, particularly if family relationships are unusual.

Other professionals (e.g. educational psychologists, therapists) may have helpful ideas to help the student learn. Families may also have ideas they would be happy to share, and may like to hear about what strategies work in the classroom. It will help the student if their family and the school work together.


Students with FASD need sensitive, empathetic staff who will personalise their learning to take account of their difficulties, which may affect sensory functioning, gross and fine motor skills, visual-motor, visual-spatial or visual-perceptual skills, attention and processing speed, expressive and receptive language; auditory and visual learning and memory, executive functioning, all of which are likely to impact on academic achievement. So, it is particularly important to use the students' strengths and interests in order to build self esteem and emotional resilience, and to improve long term outcomes across the curriculum and throughout life.

Blackburn (2010; Blackburn et al., 2010), through NOFAS-UK, the UK National Organisation on Fetal Alcohol Syndrome, has recently developed UK guidance and frameworks for the education of children and young people with FASD in association with schools which can be found on the organisation's website (www.nofas-uk.org).

Key references

A full reference list can be found on the Information Sheet relating to this topic.

Blackburn, C. (2010) Facing the challenge and shaping the future for primary and secondary aged students with Foetal Alcohol Spectrum Disorders (FAS-eD Project). London: NOFAS-UK. [Online at:



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

www.nofas-uk.org; accessed: 2.3.11]

Blackburn, C., Carpenter, B. and Egerton, J. (2010) Shaping the future for children with foetal alcohol spectrum disorders. *Support for Learning*, 25 (3), 139-145.


British Medical Association (2007) *Fetal Alcohol Spectrum Disorders: A guide for healthcare professionals*. London: British Medical Association.

Clarren, S (2004) *Teaching Students with FASD*. Alberta, Canada: Alberta Learning.

Connor, P D and Huggins, J (2005) Prenatal development: fetal alcohol spectrum disorders. In: K. Thies (ed) *Handbook of Human Development for Healthcare Professionals*. Sudbury, MA: Jones and Bartlett

Fetal Alcohol Syndrome Consultation, Education and Training Services, Inc. Publishers. [Online at: <http://books.google.co.uk/books?id=CkbMiPxwvBQC>; accessed: 20.4.09]

Ward, L, Mallett, R, Heslop, P and Simons, K (2003a) Transition planning: how well does it work for young people with learning disabilities and their families?, *British Journal of Special Education*, 30 (3), 132-137.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

What is foetal alcohol spectrum disorder?

The term foetal alcohol spectrum disorder (FASD) is an umbrella term used to encompass the range of possible effects of prenatal exposure to alcohol (British Medical Association, 2007), including foetal alcohol syndrome (FAS), partial foetal alcohol syndrome (pFAS), foetal alcohol effects (FAE), alcohol related neurodevelopmental disorder (ARND) and alcohol related birth defects (ARBD). Any amount of alcohol consumed by a mother during pregnancy crosses the placenta, and can result in birth defects for her child, including physical, mental, behavioural and/or learning disabilities, with life long implications. Full FAS is associated with characteristic physical effects including facial dysmorphism, but, even where there are no observable effects, children are left with irreversible brain damage.


Foetal alcohol exposure is the leading known cause of intellectual disability in the Western world, and according to international studies, it is estimated that one in every 100 children is born with FASD (Autti-Ramo, 2002; British Medical Association, 2007; May and Gossage, 2001; Plant, 1985; Plant et al, 1999; Sampson et al, 1997). This is greater than the combined incidence of children born in any year with Down syndrome, cerebral palsy, cystic fibrosis and spina bifida. In a culture which sees binge drinking on the increase (Donaldson, 2009), the number of children with FASD is set to rise. Based upon the above estimate in the context of annual birth figures produced by the Office of National Statistics, there are likely to be 6–7,000 babies born with FASDs of varying severity in the UK each year.

Teachers and teaching support staff will undoubtedly meet students with FASD in their classrooms. They need to know how to respond to their learning needs effectively, enable them to maximise their potential, improve their life chances, and take their places alongside their mainstream peers as citizens (DfES, 2004; HM Government, 2004). To do this, teaching staff will need training and support to realise this in the context of the English National Curriculum and National Education Strategies (cf. <http://nationalstrategies.standards.dcsf.gov.uk/>) (Carpenter, 2009; 2011). In other countries (eg Canada, USA), research outcomes have led to improved educational support for students with FASD.

Four criteria, including growth deficiency, specific facial features, central nervous system damage and confirmed prenatal alcohol exposure must all be met for a full diagnosis of FAS. However, many students not meeting the full diagnostic criteria may still be affected and experience difficulties. For example:

- ARBD includes characteristics such as heart defects, sight/hearing problems, joint defects, etc
- ARND and FAE include attention deficits, behaviour disorders, obsessive/compulsive disorder.

The severity of presentation (eg facial dysmorphism) is not necessarily indicative of the severity of impairment (Stratton et al, 1996), and therefore it is also important that teachers are aware of the true effects of the hidden impairments associated with FASD, so they can recognise and accommodate students' learning needs. These students may score within normal limits on measures of IQ, and give the appearance of functioning at a level consistent with their chronological age. They appear physically mature, and are able to meet basic literacy demands and to use relatively sophisticated language. However, their academic abilities are below their IQ level, and their living skills, communication skills and adaptive behaviour levels are even more so (Streissguth et al, 1996). A lack of awareness of the difficulties of students' difficulties can lead to consistently unrealistic expectations. Without the



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

appropriate supports and interventions, this can cause them to develop serious behavioural, cognitive, and psychological secondary disabilities.

Fetal Alcohol Syndrome Consultation, Education and Training Services, Inc.
(<http://www.fascets.org/info.html>) describe the learning issues for students with FASD as follows:

Primary characteristics of individuals with FASD

No one or two characteristics are necessarily diagnostically significant; many overlap with those of other diagnoses, eg ADD/ADHD, learning disabilities, and others


- memory problems
- difficulty storing and retrieving information
- inconsistent performance (on and off) days
- impulsivity, distractibility, disorganisation
- ability to repeat instructions, but inability to put them into action ('can talk the talk but can't walk the walk')
- difficulty with abstractions, such as maths, money management, time concepts
- cognitive processing deficits (may think more slowly)
- slow auditory pace (may only understand every third word of normally paced conversation)
- developmental lags (may act younger than chronological age)
- inability to predict outcomes or understand consequences

Primary characteristics of individuals with FASD

- highly verbal
- bright in some areas
- artistic, musical, mechanical
- athletic
- friendly, outgoing, affectionate
- determined, persistent
- willing
- helpful
- generous
- good with younger students

Secondary characteristics which individuals with FASD may develop due to chronic frustration

If their condition goes unrecognised, and they do not receive support, patterns of defensive behaviours commonly develop over time. These characteristics are believed to be preventable with appropriate supports.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

- fatigue, tantrums
- irritability, frustration, anger, aggression
- fear, anxiety, avoidance, withdrawal
- shut down, lying, running away
- trouble at home, school, and community
- legal trouble
- drug / alcohol abuse
- mental health problems (depression, self injury, suicidal tendencies)

The importance of providing appropriate support for students with FASD cannot be emphasised enough. The secondary behaviours described above may become disabling. Research describes the bleak outcomes for some young people with FASD: mental health problems (seen in 87% of children; O'Connor et al, 2002); disrupted school experience (60% over the age of 11 years; Riley, 2003); trouble with the law (60% of teenagers; Kelly, 2009); imprisonment (50%; Kelly, 2009); inappropriate sexual behaviour; problems with dependent living (80%; Riley, 2003) and employment (Streissguth and Kanter, 1997). They also are at increased risk of developing addictive behaviours such as alcohol abuse, thereby potentially continuing the cycle of FASD into the next generation (Baer et al, 2003). Streissguth and colleagues (1996) found that 3% of 6–11-year-olds, 12% of 12–20-year-olds, and 23% of adults from a cohort of 415 subjects diagnosed with FAS or Foetal Alcohol Effects had attempted suicide. (The adult figure is five times the US national average.)

Teaching and learning

The need for personalised, meaningful and high quality education is crucial if we are to divert this outcome. In developing personalised learning pathways for students with FASD, practitioners need to take account of their levels of impairment in terms of: sensory perceptual functioning; gross and fine motor skills; visual-motor integrative abilities; visual-spatial and visual-perceptual skills; attention and processing speed; expressive and receptive language; auditory and visual learning and memory; executive functioning; IQ and academic abilities.

It is important to build upon the positive personality characteristics, strengths and talents of students with FASD (Alberta Learning, 2004), and to manage their learning environment to allow them to flourish. This may include providing consistency, structure and repetition, sensory regulation, and a concrete, hands-on approach to learning. Students with FASD often have strong visual memories and good verbal fluency. They often have high energy levels, and a gregarious, fun loving, caring and affectionate nature. Many are skilled in visual arts and music, and different sports.

The adapted SCORES approach (Clarren, 2004; Lasser, 1999; see table below) summarises effective approaches for working with students with FASD. The summary emphasises the importance of structure. However, wherever possible, it is important that this structure is visual. A structured teaching approach, such as TEACCH (Mesibov et al, 2006), provides specific guidance about how this can be done. This may include:



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

- creating clearly defined areas of the classroom which are used consistently for specific activities (eg a relaxation area, an area for independent work, etc)
- using screens or similar to reduce the visual distraction around the student's work area
- using visual timetables and visual systems (eg written, symbolled, pictorial, etc) to show the order of different tasks or the order of elements within a task; the level of complexity and communication should be tailored to the individual student
- organising work to be carried out in a systematic way using labelled shelves, trays, etc.

SCORES teaching and learning approaches (Clarren, 2004; Lasser, 1999)

S

Supervision

- Supervise students closely to keep them safe and prevent problems.

Structure

- Teach students that every day has a consistent structure to it.
- Make sure routines are explicit, firmly in place and followed.
- Ensure each class and every learning activity is planned and structured.
- Use task analysis to ensure that all steps required to complete an assignment are given and understood.
- Make sure directions are simple, and given orally and in visual form.

Simplicity

- Keep everything simple – rules, routines, directions, language, explanations and expectations.

Support

- Provide unconditional emotional support to the student.
- Ensure support for students' families and teachers as necessary to deal with emotional issues such as grief, loss and frustration.

Success


- Identify students' strengths and help them recognise and use their own strengths.
- Look for positive events, set up situations to ensure accomplishments and celebrate success.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

C	<p>Communication</p> <ul style="list-style-type: none"> • This takes place regularly between the home and school. • Everyone involved with these students knows what others are doing to help and communicates when there are changes in plans. • Teach students how to communicate feelings and needs. <p>Consistency</p> <ul style="list-style-type: none"> • Routines, rules and expectations need to be consistent. • Give steps to complete a task in the same way every time.
O	<p>Organisation</p> <ul style="list-style-type: none"> • Teach organisation skills in the classroom. • The classroom is organised—a place for everything and everything in its place. • Learning activities and daily routines are organised.
R	<p>Rules</p> <ul style="list-style-type: none"> • They are simple and easy to follow. • Use concrete and positive language — 'Walk' rather than negative 'Don't run' or abstract 'Be careful'. • All staff use the same words for each rule. • Check whether students know and understand what the rules mean. • If a student does not follow a rule, an adult corrects the behaviour immediately, without scolding, and encourages the student to try the behaviour again, this time following the rule.
E	<p>Expectations</p> <ul style="list-style-type: none"> • Expectations need to be realistic, attainable and easily understood. • Take into consideration special and individual needs of students for life and social skills as well as academics. • Clearly specify what is to be expected and accomplished for any given task or activity.
S	<p>Self-worth</p> <ul style="list-style-type: none"> • Students feel accepted, valued and safe. • Give positive encouragement each day. <p>Build on students' strengths to help them cope with the frustration of things they cannot do.</p>



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

Current research indicates that three main areas offer the main challenge to teachers (Yukon Education, 2006; Kleinfeld and Wescott, 1993):

Numeracy and mathematics

Researchers in neuroscience have reported that in many children with FASD, the parietal lobe is severely affected (Goswami, 2004; Kopera-Frye et al, 1996). This area of the brain deals with numeracy and mathematical computation. The teaching challenge here is more than a straightforward differentiation of the mathematics curriculum.

Executive function and behaviours for learning

The lack of structure and self discipline means that students with FASD are often erratic and unfocused. They lack the basic organisational skills that are fundamental to effective learning. Their disorientation in the classroom environment leads them to disengage quickly from the learning flow. Cumulatively, this means that they do not make satisfactory learning gains or adequate progress in learning management.


PSHE (emotional wellbeing)

The literature in this area has widely reported the vulnerability of young people with FASD to mental health problems. Their lack of social skills, and difficulties informing sustainable friendships, makes them susceptible to feelings of negativity and poor self-esteem. US studies have reported high levels of suicide amongst young adults with FASD (Streissguth et al, 1996). We believe that targeted educational

The cognitive and behavioural profile of students with FASD changes over time, so the learning needs of primary and secondary students are subtly different. Learning, behavioural/emotional and social difficulties typically become more evident as they progress through school. Therefore, repeated neuropsychological assessment may be needed at different times during the life of an individual with FASD to capture their evolving strengths and weaknesses accurately, and to plan appropriate interventions.

Transition between primary and secondary schools needs to be carefully managed, as this is an area in which support strategies and services can often become disrupted, and communication can break down between practitioners (Ward et al, 2003). For teenagers, issues around emotions, friendships and sexual behaviour, independence and achievement compound their primary impairments (Connor and Huggins, 2005).

Whatever the background, the challenge remains, 'How do we optimise learning for this student group?'. There is as yet no direct guidance from any government agency in the UK to teachers on how to educate students with FASD. However, currently the Training and Development Agency for Schools (TDA) are supporting the FAS-eD research project which is looking into effective teaching and learning approaches for children with FASD (reporting October 2010). There is also information available on working with children with FASD in the early years from Worcestershire County Council/Sunfield School (Blackburn, 2009).




COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

Key references

- Alberta Learning (2004) *Teaching Students with Fetal Alcohol Spectrum Disorder: Building strengths, creating hope*. Edmonton, Canada: Alberta Learning.
- Autti-Ramo, I (2002) Foetal Alcohol Syndrome: a multifaceted condition, *Developmental Medicine & Child Neurology*, 44, 141–144.
- Benton Gibbard, W, Wass, P and Clarke, M E (2003) The neuropsychological implications of prenatal alcohol exposure, *Canadian Child and Adolescent Psychiatry Review*, 12 (3), 72–76.
- Blackburn, C. (2010) Facing the challenge and shaping the future for primary and secondary aged students with Foetal Alcohol Spectrum Disorders (FAS-eD Project). London: NOFAS-UK. [Online at: www.nofas-uk.org; accessed: 2.3.11]
- Blackburn, C., Carpenter, B. and Egerton, J. (2010) Shaping the future for children with foetal alcohol spectrum disorders. *Support for Learning*, 25 (3), 139–145.
- British Medical Association (2007) *Fetal Alcohol Spectrum Disorders: A guide for healthcare professionals*. London: British Medical Association.
- Carpenter, B (2005) Early childhood intervention: possibilities and prospects for professionals, families and children, *British Journal of Special Education*, 32 (4), 176–183.
- Carpenter, B. (2009; 2011) Pedagogically bereft: Improving learning outcomes for children with Foetal Alcohol Spectrum Disorders. *British Journal of Special Education*.
- Clarren, S (2004) *Teaching Students with FASD*. Alberta, Canada: Alberta Learning.
- Connor, P D and Huggins, J (2005) Prenatal development: fetal alcohol spectrum disorders. In: K. Thies (ed) *Handbook of Human Development for Healthcare Professionals*. Sudbury, MA: Jones and Bartlett Publishers. [Online at: <http://books.google.co.uk/books?id=CkbMiPxwvBQC>; accessed: 20.4.09]
- Department for Children, Schools and Families (2008) *The Children's Plan: One year on*. Annesley, Notts: DCSF. [Online at: <http://www.dcsf.gov.uk/oneyearon/ae/uploads/documents/flagship.pdf>; accessed: 27.4.09]
- Department for Education and Skills (2004) Removing Barriers to Achievement: *The government's strategy for SEN*. Annesley, Notts.: DfES Publications. [Online at: <http://www.standards.dfes.gov.uk/eyfs/resources/downloads/removing-barriers.pdf>; accessed: 20.4.09]
- Donaldson, L (2009) *Annual Report of the Chief Medical Officer*. London: Department of Health. [Online at: http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/AnnualReports/DH_096206; accessed: 23.8.10]



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

Florida Department of Education (2005) *Teaching Students with Fetal Alcohol Spectrum Disorders*.
Florida: Florida State University Center.

Goswami, U. (2004) Neuroscience, education and special education, *British Journal of Special Education*, 31 (4), 175–183.

Green, C R, Mihic, A M, Brien, D C, Armstrong, I T, Nikkel, S M, Stade, B C, Rasmussen, C, Munoz, D. P and Reynolds, J N (2009) Oculomotor control in children with fetal alcohol spectrum disorders assessed using a mobile eye-tracking laboratory, *European Journal of Neuroscience*, 29 (6), 1302–1309.

HM Government (2004) *Every Child Matters: Change for children*. [Online at: <http://www.everychildmatters.gov.uk/>; accessed: 20.4.09]

Huggins, J E, Grant, T, O'Malley, K and Streissguth, A P (2008) Suicide attempts among adults with fetal alcohol spectrum disorders: clinical considerations, *Mental Health Aspects of Developmental Disabilities*, April–June. [Online at: http://findarticles.com/p/articles/mi_6883/is_2_11/ai_n28524972/?tag=content;col1; accessed: 27.4.09]

Jones, K L and Smith, D W (1973) Recognition of the fetal alcohol syndrome in early infancy, *Lancet*, 2 (7836), 999–1001.

Kelly, K (20.4.09) Is foetal alcohol spectrum disorder linked to anti-social behaviour?. Woman's Hour, Radio 4, 10–11 a.m. [Online at: http://www.bbc.co.uk/radio4/womanshour/03/2009_16_mon.shtml; accessed: 20.4.09]


Kemmis, S and Wilkinson, M (1998) Participatory action research and the study of practice. In: B Atweh, S Kemmis and P Weeks (eds) *Action Research in Practice: Partnership for social justice in education*. London: Routledge.

Kodituwakku, P, Coriale, G, Fiorentino, D, Arago'n, A S, Kalberg, W O, Buckley, D, Gossage, J P, Ceccanti, M and May, P A (2006) Neurobehavioral characteristics of children with fetal alcohol spectrum disorders in communities from Italy: preliminary results, *Alcoholism: Clinical experimental research*, 30 (9), 1551–1561.

Kopera-Frye, K, Dehaene, S and Streissguth, A P (1996) Impairments of number probably induced by prenatal alcohol exposure, *Neuropsychologia*, 34, 1187–1196.

Lemoine, P, Haroussseau, H, Borteyru, J P, et al. (1968) Les enfants de parents alcooliques: anomalies observées, à propos de 127 cas, *Ouest Médical*, 21, 476–482.

Lemoine, P and Lemoine, P (1992) Avenir des enfants des mères alcooliques (études des 105 cases retrouvés à l'âge adulte) et quelques constations d'intérêts prophylactiques, *Annales de Pédiatrie*, 39, 226–235.



COMPLEX LEARNING DIFFICULTIES AND DISABILITIES RESEARCH PROJECT (CLDD)

FOETAL ALCOHOL SPECTRUM DISORDERS

May, P A and Gossage, J P (2001) Estimating the prevalence of fetal alcohol syndrome: a summary. [Online at: <http://pubs.niaaa.nih.gov/publications/arh25-3/159-167.htm>; accessed: 19.4.09]

Mukherjee, R A S, Hollins, S and Turk, J (2006) Fetal alcohol spectrum disorder: an overview, *Journal of the Royal Society of Medicine*, 99, 298–302. [Online at: http://www.intellectualdisability.info/mental_phys_health/fetal_alcohol_mukherjee.htm; accessed: 19.4.09]

National Children's Bureau (NCB) (2003) *Guidelines for Research*. [Online at: http://www.ncb.org.uk/Page.asp?originx1838gs_81330095622449n31q531441; accessed: 23.4.09]

O'Connor, M J, Shah, B, Whaley, S, Cronin, P, Gunderson, B and Graham, J (2002) Psychiatric illness in a clinical sample of children with prenatal alcohol exposure, *American Journal of Drug and Alcohol Abuse*, 28, 743–754.

Plant, M L (1985) *Women, Drinking and Pregnancy*. London: Tavistock.

Plant, M L, Abel, E and Guerri, C (1999) Alcohol and pregnancy. In: Macdonald, I (ed) *Health Issues Related to Alcohol Consumption* (2nd edn). Oxford: Blackwell, 181–214.

Sampson, P D, Streissguth, A P, Bookstein, F L, Little, R E, Clarren, S K, Dehaene, P, Hanson, J W and Graham, J M Jr (1997) Incidence of fetal alcohol syndrome and prevalence of alcohol-related neurodevelopmental disorder, *Teratology*, 56 (5), 317–326.

Streissguth, A, Barr, H, Kogan, J and Bookstein, F (1996) *Understanding the Occurrence of Secondary Disabilities in Clients with Fetal Alcohol Syndrome (FAS) and Fetal Alcohol Effects (FAE. Final Report: Centers for Disease Control and Prevention Grant No. 04/CCR008515)*. Seattle, WA: University of Washington Fetal Alcohol and Drug Unit.

Streissguth, A and Kanter, J (eds) (1997) *The Challenge of Fetal Alcohol Syndrome: Overcoming secondary disabilities*. Washington, DC: University of Washington Press.

Streissguth, A P and O'Malley, K (2000) Neuropsychiatric implications and long-term consequences of fetal alcohol spectrum disorders, *Seminars in Clinical Neuropsychiatry*, 5 (3), 177–190.

Visser, J (2009) *Diversity and Personalised Learning*. London: Routledge.

Ward, L, Mallett, R, Heslop, P and Simons, K (2003a) Transition planning: how well does it work for young people with learning disabilities and their families?, *British Journal of Special Education*, 30 (3), 132–137.

